

Community Forest Management: The story behind a success story in Nepal

François Libois, Jean-Marie Baland, Nicolas Delbart,
Subhrendu Pattanayak

December 20, 2021

DeFiPP Working Paper 2021-06

Community Forest Management: The story behind a success story in Nepal

François Libois*, Jean-Marie Baland†, Nicolas Delbart‡, Subhrendu Pattanayak§

December 20, 2021

Abstract

Since 1993, Nepal implemented one of the most ambitious and comprehensive program of decentralization of forest management in the world, widely considered as a success story in terms of participatory management of natural resources. Using quasi-experimental methods, we first quantify the net gains in tree cover related to the program in the Hills and Mountains of Nepal and describe their temporal evolution. We then discuss the mechanisms driving forest restoration, highlighting that while community forestry played a role in increasing forest biomass and forest size, it also reduced demand pressures by altering energy choices.

Keywords: Forest management ; Community forestry ; Nepal ; Energy ;
Participatory development

JEL codes: O13, Q23, Q5, D23, D12

*Paris School of Economics and INRAE, France. Email: francois(dot)libois(at)psemail(dot)eu

†CREDES, University of Namur, Belgium, BREAD and CEPR. Email: jean-marie(dot)baland(at)unamur(dot)be

‡LIED, University of Paris, France. Email: nicolas(dot)delbart(at)u-paris(dot)fr

§Duke University. Email: subhrendu(dot)pattanayak(at)duke(dot)edu

1 Introduction

Over the past 25 years, the government of Nepal implemented one of the most ambitious and comprehensive program of decentralization of forest management in the world. This major institutional change resulted in the transfer of the management of almost 50% of Nepal forests to no less than 18000 Community Forest User Groups (CFUG). More than a third of the Nepalese population are directly involved in the management of forests, a key natural resource in everyday life which provides not only firewood or timber, but also fodder for the livestock, fruits, nuts or medicinal plants. Thus, in the Hills and the Mountains, about 45% of rural households report their first source of firewood as being the community forest.

The program, formally launched in 1993, entrusts forests to Community Forest User Groups at the village level, who are then in charge of their daily management. Through their formal constitution and their operational plans, they lay down the rules of access and use of the forest, manage their monitoring and develop planting and harvesting programs. User fees and the sales of timber and other forest products generate income which is reinvested in the forest or used for collective projects and public infrastructure at the village level. These income sources are important, and largely exceed the budget of the 'local development committee'. (In a private conversation, the Head of the CFUG Division in the Forest Department estimated that CFUGs incomes were four times larger than the total budget of the local village committees.) The success of the program has been widely advertised and received lot of attention internationally, for instance by UNEP (Sukhdev and Nuttall, 2010).

However, we still know very little about the effects of the program at the country level and about the potential channels underlying these positive changes (For a similar assessment at the world scale, see Bowler et al. (2012)). From recent studies in Nepal, the evidence is mixed. Using propensity score matching on a broad sample of community forests in Nepal, Bluffstone et al. (2018) finds that formal CFUGs do not sequester more carbon than forests under informal community management. Oldekop et al. (2019) compares changes in forest cover and poverty following the creation of community forest user groups between 2001 and 2011. They find that subdistrict that are otherwise similar at baseline tend

to experience reductions in deforestation and in poverty. At a more limited scale (in the Arun Valley at early stages of the program), Edmonds (2002) finds that the creation of CFUGs reduces firewood collections at the household level.

In the neighbouring Indian Himalayas, Somanathan et al. (2009) compares forests under community management (Van Panchayat) to neighbouring forests, using satellite imagery. They conclude that, compared to situations of open and unregulated access, Van Panchayats are as effective in preserving forest cover in community forests than the Forest Department in State forests, but at a much lower cost in terms of fiscal resources and bureaucratic management. Using a cross section of forest measures taken in randomly chosen forest patches, Baland et al. (2010) show that, despite the fact Van Panchayats are initially given more degraded forests, they rapidly succeed in reducing excessive lopping and tree damages, leading to a much healthier and denser forest in the long run. By contrast, in the context of Madagascar during economic and political crises of the early 2000's, Desbureaux (2016) shows that community forest management led to increased deforestation, particularly in villages neglected by the central authorities in which local collective action was traditionally strong. With the possible exception of the civil war that ended in 2006, the situation in Nepal differs from the latter as the Forest Department in Nepal strongly supported and accompanied the community forest policy in the creation of Forest User Groups in the villages.

In this paper, we first show that the CFUG program contributed to substantial increases in forest cover in the Hills and the Mountains of Nepal. To this end, we create a 14 years panel data set that combines remote sensing data with administrative data and household surveys. Given that the creation of CFUGs cannot be considered as fully random, we model the spread of the program in a district by instrumenting the creation of a CFUG with the interaction between the time since the start of the program in a given district and the distance between a given village and the district headquarters. This approach allows to isolate the effect of CFUGs on forest cover, independently of the fact that, for instance, CFUGs may have been created in more degraded or less valuable forests. We also investigate the time structure of these effects, which reveals a steady increase in canopy density, consistent with an immediate reduction in lopping, followed by a slow and steady regeneration process. We then investigate the potential mechanisms

underlying these forest improvements. On the supply side, forest areas increased substantially at the expense of agricultural land and shrubs, and replanting took place, with needle tree and mixed forests increasing much more than broadleaf forests. CFUGs also reduce the demand for biomass. We find that small-scale biogas installations, a direct substitute to firewood for cooking, are widely adopted in areas of CFUGs expansion. Using household data, we also find that firewood collection times are higher and firewood collection lower in villages in which CFUGs are recent, while these correlations vanish for older CFUGs. This evidence is consistent with the idea that CFUGs start first by imposing restrictions on firewood collections but later allow, as forests conditions improve, larger collections of forest products, and in particular firewood, by the villagers.

In the following, we leave the Terai region aside as the Terai is specific in several ways. First, forests have long been cleared in Terai, with remaining patches of forests in the Northern part of the Terai in the first slopes of the Siwalik. Second, Terai forests are mostly covered by sal trees (*Shorea robusta*), a highly valued commercial species traded on the legal and illegal markets, particularly along the Indian border. These two features completely change the nature of local community management there (see Libois (2021)). Third, because of the milder climate in those plains, energy needs do differ, particularly for heating in the winter season. Finally, the methodology adopted in this paper is much less appropriate for the Terai forests.¹

The paper is organized as follows. Section 2 describes the data used. In section 3, we present the effects of community forestry on forest conditions. Section 4 discusses potential mechanisms behind the gains in tree cover while the last section concludes.

¹First, the Leaf Area Index is less precise in Terai as the seasonality of cropping patterns is much less standardized, with green fields persisting in November and December. Second, the dynamics of CFUG creation in the Terai differs as transportation costs across these flat plains are much lower.

2 Data

To measure tree cover, we rely on two main data sources. First, we use the yearly village-average November leaf area index (LAI) constructed using the algorithm by Verger et al. (2014) applied to the data from the SPOT-VEGETATION sensor over the years 1999 to 2013 and to the data from the PROBA-V sensor from 2014 onwards.² LAI is a vegetation parameter commonly used to monitor the spatial and temporal variation in the leaf density. Considered as one of the Essential Climate Variable (Bojinski et al., 2014), it is defined as half of the area covered by all the leaves per ground unit area (Bréda, 2008). The leaf area index gives us a continuous measure of tree cover that reacts to increases in leaf density. (It is less subject to saturation than other possible measures, thereby allowing to better capture variations in canopy density.) We average the LAI at the 'village' level where the 'village' corresponds to the smallest administrative division in Nepal, corresponding to the Village Development Committee (VDC), often composed of different hamlets. Henceforth, we refer to a VDC as a 'village'. We use November data because, in the Hills and Mountains of Nepal, the deciduous trees still have their green leaves whereas the cultures have just been sown. In that period, the green photosynthesis visible by remote sensing comes essentially from tree leaves (Niraula et al., 2013). This observation has been used in several remote sensing based methods to map forest cover in Nepal (ICIMOD, 2014a,b), and has been corroborated by our field visits and observations of the phenology of the different types of vegetation.

We also rely on the MODIS land cover type product (MCD12Q1), a data set that provides an annual classification of land cover at 500m resolution. The product is created using supervised classification of MODIS reflectance data (Friedl et al., 2002, 2010). We use the class description of the International Geosphere-Biosphere Programme and focus our analysis on forest related classes (see Sulla-Menashe and Friedl (2018) for more details). We also make extensive use of the 2017 census of CFUG provided by the Department of Forest in Nepal, which

²The dataset relies on a neural-network based algorithm trained with older generation LAI datasets in order to merge their respective pros. It also includes a procedure in order to make up for the gaps in the time series due to the presence of clouds, as cloud cover can occur even outside the rainy season (June-September).

records the creation date of all groups since the inception of the program. It also contains information about the area managed by the CFUG, its administrative location as well as some information about the membership of the group and the steering committee composition.³ Our main variable of interest is the share of a village area managed by the CFUG, the village area being measured with a digitized map of the village boundaries established by the Nepal Central Bureau of statistics.

In figure 1 we plot the evolution of the November Leaf Area Index and the share of village area managed by CFUG over the last 20 years. The overall trend in tree cover is overall positive and correlated with a sharp increase in the area managed by CFUG, with one eighth of the territory under community management at the end of the period.

[Insert Figure 1 here]

2.1 Additional data sources

We also make use of additional data sources to investigate the mechanisms driving the evolution of tree cover. We first exploit the census of biogas installations of Nepal over the period 1994-2015 from the Alternative Energy Promotion Center. It contains very detailed information on all biogas installations subsidized by the Nepal government and constitutes the most comprehensive database on biogas at the village level in Nepal.

Our main source of information on household choices are the second and third wave of the World Bank Living Standards Measurement Survey (LSMS) for Nepal, also known as the Nepal Living Standard Survey (NLSS). The Nepal Central Bureau of Statistics, in collaboration with the World Bank, interviewed households about several aspects of their production and consumption activities. The surveys cover 123 villages and 1474 households in 2003-4 and 178 villages and 2116 households in 2010-11 in the Hills and Mountains region, selected randomly with a probability proportional to their population. The quality of the surveys has been tested by Hatlebakk (2007), who also discusses them in greater details. CBS

³We also use the 2010 CFUG census once to provide descriptive statistics about the type of forests managed by CFUG.

(2011) provides additional information about the technicalities of the sampling, the methodology, and the implementation of the surveys.

Our analysis relies extensively on additional controls derived from various data sources. To compute walking distances between district headquarters and villages, we use the third version of the 30m-resolution ASTER digital elevation model (Abrams and Crippen, 2011; Fujisada et al., 2011).⁴ The same source of information provides village level median elevation and its standard deviation. Temperature related controls were extracted from the MODIS product MOD11A2, that provides 8-day average land surface temperature at a 1km-resolution (Wan et al., 2015). Snow cover comes from MOD10A2 product, a 8-day snow cover measure provided by Hall and Riggs (2021) at a 500m-resolution. Lastly, we compute village level annual rainfall based on the daily estimates of the tropical rainfall measurement mission (TRMM, 2011).

While remote sensing allows to have high frequency information on the environment, data become less frequent and less precise as we go back in time. We have therefore digitized and geocoded US army paper maps of Nepal from the 1950s. Using a semi-automatic classification tool, we have extracted areas depicted as forest on these maps.⁵ This information gives us a historical measure of forest cover around 1950, largely before the start of the community forestry program.⁶ Lastly, the Informal Sector Service Centre (INSEC), a very active Nepali human rights organization, collected extensive data on conflict intensity during the whole duration of the civil conflict. The INSEC database (INSEC, 2009) is considered as the most reliable data source on the civil war that lasted over the period 1996-2006. For more details on this data source see Libois (2016) and Joshi and Pyakurel (2015). Table (1) and (2) below provide the main descriptive statistics of the village and household variables used throughout our estimations. Some additional household information is given in Table A1 in the Appendix.

[Insert Table 1 here]

⁴We follow the formula of Aitken (1977); Langmuir (2013) based on Naismith’s rule of thumb to compute walking time and implemented in GRASS GIS

⁵We could rely on the Historical Map plugin in QGIS to perform this analysis.

⁶We use this variable with caution as it pre-dates huge population expansion in Nepal as well as the colonization of Terai. This internal migration from the Hills to the plains was accompanied by a huge deforestation process in the low lands as well as structural changes in the Hills.

[Insert Table 2 here]

3 Community forestry and forests

3.1 Empirical strategy

To investigate the consequences of the community forestry program, we first compare forest conditions, as measured by tree cover at the village level, before and after the creation of a community forest user group. Given the dynamics of forest regeneration, we then analyse the evolution of tree cover over time, once a forest user group is created.

The main empirical strategy relies on the following specification:

$$LAI_{vt} = \alpha CFUGshare_{vt-1} + \beta_k X_{kvt} + \eta_v + \delta_t + \varepsilon_{vt} \quad (1)$$

where LAI , stands for the average leaf area index in village v in year t . The main explanatory variable, $CFUGshare$, is the share of the village area under CFUG management in year t . In all specifications, we control for a set of time-varying village level controls X such as total rainfall, average snow cover, growing degree days or the local intensity of the civil conflict. We also include village and time fixed effects parameters, to avoid potential biases caused by village characteristics or country level shocks, such as altitude or national political cycles. The parameter of interest, α , quantifies the change in the average leaf area index that follows a change in the share of the area managed by the village CFUGs.

As such, a causal interpretation of α is questionable, given the non-random nature of the creation of CFUG. For instance, if members of CFUG are systematically more pro-social, with stronger social ties, and more aware of environmental issues, we may expect increases in LAI to occur in places where CFUGs are created, independently of the intrinsic properties of the program, biasing upwards the estimated α . Alternatively, the latter may be downward biased if the forest department systematically chooses to hand over the less productive or more degraded forest plots to local communities. To reduce these potential biases, we follow an instrumental approach and estimate equation (1) by two-stage least squares. The

first stage equation is given by:

$$CFUGshare_{vt} = \beta_1 Proximity_v \times TO_{dt} + \mathbf{Z}_{vt} \Theta + \gamma_v + \tau_t + \varepsilon_{vt} \quad (2)$$

The instrument used is the interaction between *Proximity*, the inverse of the distance between the village the district headquarters where the district forest office is located (as measured by the walking time in hours), and *TO*, the number of years since the onset of the CFUG program in district *d*. We again control for a large set of time and space varying variables, including *TO* and a broad range of environmental controls, as well as village and time fixed effects. The instrument we propose is relevant both from a statistical point of view and given our insights from the field and interviews with forest officers. The first CFUGs were typically created close to district forest offices after the nomination of a forest officer willing to implement this new program. These creations initially involved numerous visits and extensive efforts to persuade villagers to join this new program. Once the program starts in a district, the first CFUGs are almost always created close to the district office of the department of forest. Indeed, in the nineties, travelling within districts would typically occur by foot or riding horses.⁷ Forest officers had therefore strong incentives to minimize travel distance.

We proxy travel time to villages by computing the fastest walking time between district headquarters and the village development committee in this district relying on a digital elevation model of Nepal.⁸ There are large variations in the launching of the CFUG program across districts, as some started in the early nineties while the last district to do so was Bara, where a first CFUG was created in 1999.⁹ We do not consider these variations as exogenous as such but assume that the interaction between the two sources of variations is an exogenous predictor of the creation of a CFUG, conditionally on the controls included in the regression.¹⁰ In a

⁷Some districts were not even connected by paved road to the capital city Kathmandu.

⁸Travel distance may be correlated with other factors influencing forest cover, which we control for by village fixed effects.

⁹There is no CFUG in Mustang because CFUG are not created in Conservation areas and the whole district is part of the Annapurna Conservation Area.

¹⁰In practice, the onset of a program in a given district has arguably some random component as it is partially driven by mutations and promotions in the department of forests, which exhibits a relatively high level of turnover.

sense, this instrument is a generalization of Edmonds (2002) approach over a larger spatial and temporal coverage. As we see in Figure 2, villages located further away from district headquarters are increasingly incorporated in the program. Figures A2, A3 and A4 in the Appendix illustrate this expansion process that looks like an oil spill starting in each district headquarters at different points in time.¹¹

[Insert Table 2 here]

On top of being statistically strong, the instrument has to be exogenous, conditionally on the other control variables. To violate this exogeneity assumption, one would have to find a variable which affects tree cover and is correlated with a district specific expansion starting from the district headquarters in the district inception year of the CFUG program. The development of infrastructure could be such a threat, but it requires the onset of the road construction programs to be correlated with the start of a CFUG program in the district. This is not what we observe in the spatial distribution of the launching of the CFUG program. Another possible threat is that economic development accelerates in a district at the same moment as the onset of the CFUG program and that economic development induces rural exodus or changes in domestic energy choices that occur at the same temporal and spatial pace than CFUG creation. This looks in our view rather unlikely.

3.2 Main result

Table 3 reports the estimation results of the effect of CFUGs on the November Leaf Area Index following equation 1. (In all our estimations, we weight the observations by the village area to reflect changes in tree cover at the country level and we cluster standard errors at the district level. This is the natural level since our instrument assumes that there are common shocks at the district level inducing CFUG creations.) Column (1) indicates that a 10% increase in the share of village area managed by a CFUG is followed by an increase in the leaf area index

¹¹In the Terai plains - the twenty Southern districts bordering India - this expansion is less systematic as it is much easier to travel across these flat areas and forests have been cleared in most places but the national parks and the foothills of the Siwalik.

of 0.04. In terms of magnitude, with an average leaf area index of 1.3 and 12% of the village area managed by a CFUG in 2013, the contribution of the CFUG program to the increase in tree cover is estimated to be about 4 %.¹²

[Insert Table 3 here]

Given our previous discussions, we expect some selection in the forest plots that the department of forest hands over to local communities. We therefore estimate the main equation using our instrument and report the results in column (3) of table 3. The two-stage least-squares approach reinforces the conclusions of the simple panel approach, and indicate larger effect of CFUGs on the leaf area index. Point estimates actually increase tenfold, and a 12% increase in CFUG coverage increases the leaf area index by 0.55, a 40% increase. These large estimates are consistent with our fields observations whereby the forest department tends to hand over forests plots that are already degraded. CFUGs are first created close to urban centres where the pressure on land is the largest and the need for forest products remain high. Forests that are easier to protect and less at risk of degradation tend to remain longer in the hands of the Department of Forest.¹³ This is a typical source of downward bias for the OLS coefficient which explains why instrumenting for CFUG creation amplifies the point estimate: given low initial levels of forest cover, changes can only be large. The first stage estimates are reported in the second column of 3. Ten years after the program starts in a district, a village located at 5 walking hours from the district headquarters is expected to have an increment of one percentage point of its area managed by CFUGs.¹⁴

In the last three columns, we report the estimations obtained when we additionally control for population density and the prevalence of biogas (in number of installations per household). One can indeed suspect that migration accelerates at the same pace as the CFUG program by starting in places close to urban centres and expanding later to more remote places. The spread of new technologies, such as biogas, could have also followed a similar pattern in time and space. These additional controls do not substantially change our results. The minor changes in the

¹² $0.04 = 0.12 \times \frac{.448}{1.33}$

¹³Baland et al. (2010) also report similar stories in the Indian Himalayas.

¹⁴ $0.011 = \frac{10}{5} \times 0.0055$. The first quartile of the walking time between a village and its district headquarters is equal to 5.57 hours.

estimated coefficients could possibly indicate that part of the CFUG effects are mediated through the adoption of biogas and the resulting reduction of firewood use. (A genuine correlation between CFUGs and biogas should have substantially reduced the estimated coefficient of CFUG). As a matter of fact, the biogas program, which involves subsidies and subcontracting to private companies, developed much later than the CFUG program, and were managed by an independent administration (the Alternative Energy Promotion Center) with no links to the Department of Forest. Our field observations indicate that biogas companies would typically take advantage of CFUG assemblies to promote their technology. The expansion of biogas can thus be viewed in some ways as a by-product of CFUG expansion, a question which we address in section 4 below. Population density should capture outmigration in the most relevant way for firewood collection, and the estimated coefficient has the expected sign. Our measure however is far from perfect, as it is based on a log linear interpolation of the information available in the 2001 and 2011 census. (Some villages are also lost due to missing data in the 2001 census.)

3.3 Longer term effects

Forest cover is a biological process and trees take time to grow. Moreover, Forest User Groups result from collective action at the village level, which also requires time to materialize. We therefore expect the effects of community forestry to be heterogeneous across time, something that could not be captured by the specification of equation (1) which provided an 'average' measure constant through time. To investigate this temporal process, we now follow an alternative approach based on the following equation:

$$LAI_{vt} = \sum_{z=0}^{20} \alpha_z \text{Proportion of VDC area managed by FUG}_{vt-z} + \mathbf{X}_{vt} \boldsymbol{\Theta} + \gamma_v + \delta_{dt} + \varepsilon_{vt} \quad (3)$$

where the leaf area index in village v at time t is a function of the share of the village area managed by a CFUG created in year $t - z$. As in the previous specifications, we control for \mathbf{X} , a vector of time varying village specific controls, γ is a vector of village fixed effects, δ captures district-time specific variations and

ε stands for the error term. The coefficients of interest, α_z , measure the change in leaf area index in year t which follows the creation of a CFUG over a given share of the village area z years before t .

We report the estimates over a time span of 20 years in Figure 3. The first years of the existence of a CFUG see a moderate increase in the leaf area index, a consequence of the first management measures which typically consist in reducing lopping, crafting rules about fodder collection and restricting firewood collection to dry wood. Over the years, we observe a gradual increase in the leaf area index consistent with a slow process of forest regeneration. After 20 years, point estimates more or less double in size compared to years closer to the creation date. The standard errors are also larger, given that fewer groups reach 20 years of age in our sample.

[Insert Figure 3 here]

In Figure 4 below, we provide an alternative estimation using the de Chaisemartin and D’Haultfoeuille (2020) approach. The latter is appropriate in our context as we have a staggered process of CFUG creation and we expect the effect of the ‘treatment’ to be heterogeneous across space and time.¹⁵ This estimator differs from the approach followed above by better defining the appropriate comparison group when estimating the ‘effect of the treatment’. It indeed compares villages which are initially similar in terms of CFUG area, but where some move to the next, higher, treatment category. Villages with lower initial shares are therefore excluded from this comparison, while they are the basis of comparison in the first approach followed above. However, this estimator requires a discrete treatment, while our measure at the village level is continuous. We have therefore decided to group CFUG coverage at the village level in six categories (one with zero, and the others with intervals of 20%). These alternative estimations show that there is no clear trend before the treatment (up to ten years) while the leaf area index increases steadily after the creation of a group.¹⁶ When compared to our

¹⁵As highlighted by de Chaisemartin and D’Haultfoeuille (2020), this pattern of treatment can generate negative weights in the OLS estimation of the average treatment effect.

¹⁶In terms of interpretation, a change of one unit in this new variable should be interpreted as a 20% increase in the share managed by CFUG.

instrumental approach in the previous sub-section, one can argue that, by properly defining the comparison group for various levels of treatment, this estimation provides an alternative way to get at the causal effect of CFUGs.

[Insert Figure 4 here]

4 Supply and demand mechanisms behind changes in tree cover

Two main reasons may explain why tree cover increases under community forest management. First, CFUG may implement better management practices, for instance by favouring some species or planting saplings. These mechanisms directly affect the supply of forest biomass. On the demand side, CFUG may reduce the demand for forest products and alleviate human pressure on forests. As shown in Baland et al. (2018), firewood collection is an important driver of forest degradation in the Hills and the Mountains. So, by modifying the energy choices of the households, CFUG may induce faster forest regeneration and tree cover gains.

4.1 The supply of biomass

We first investigate changes in land use based on the Vegetation Cover Fields of MODIS, which distinguishes between different types of vegetation covers. Table 4 follows the specification of equation 1 on a set of land use variables. The estimations indicate that a 10% increase in a village area managed by CFUG translates into an average gain of 0.8 percentage points in the overall share of forest in the village. When instrumenting for the creation of CFUG, this effect goes up to 5.8 percentage points, which is sizeable (column (5)). The large increase in forest cover is mostly driven by an increase in mixed forests and, more modestly, in needle leaf forests (columns (6) and (7)).

[Insert Table 4 here]

By contrast, as shown in Table 5, we observe no significant changes in broad-leaf forests, while the area covered with crops and shrubs decreases substantially

following the introduction of CFUGs in the village. The increase in forest cover is therefore driven by mixed and needle forests replacing crops, open (deforested) land and shrubs, which fosters overall biodiversity, particularly in mixed forests. These changes follow partly from the plantation activities undertaken by CFUGs, particularly *Pinus Roxburghii* and *Shorea robusta* for their market value as a source of timber. ¹⁷

[Insert Table 5 here]

4.2 The demand for biomass

4.2.1 Community forestry and households access to energy

Community forestry also affects the demand for energy by the villagers. By restricting access to forests and limiting firewood collections, CFUG encourage the development of alternative energy sources, such as individual biogas production units. In table 6, we show that the construction of biogas installations, whether measured by the number of biogas units in the village or the number of units per capita, increases with the presence of CFUGs in the village. In terms of magnitude, a 10% increase in CFUG coverage increases the number of biogas installations in a village by 407 units, which corresponds to an increase in terms of coverage per household of about 8.5 percentage points (column 5).

[Insert Table 6 here]

This is sizeable and should be put in perspective. First, biogas units in Nepal are rather small and require a couple of cattle heads to operate. The penetration rate of this technology is high and concerns, according to the Alternative Energy Promotion Center, 4% of the households in 2013, with an average of about 50 installations already constructed in each village of the Hills and the Mountains.

¹⁷In the 2010 CFUG census which contains information on forest type at creation date, needle forests (mostly *Pinus Roxburghii*) represent 29% and Sal forest (*Shorea robusta*) 34% of the area operated by CFUGs. *Schima Castanopsis* accounts for 10% of the CFUG forest area and Subtropical deciduous forest, 14%. *Alnus Nepalensis*, oak, rhododendron and upper slope mixed hardwood forest are more marginal. See Figure A5 in appendix for more details.

Second, according to our field visits, managers of biogas companies take advantage of CFUG assemblies to promote this technology and enter into contact with potential customers. Moreover, CFUGS themselves affect access to energy, and may offer support to biogas adoption by providing credit or subsidies.

4.2.2 Community forestry and access to firewood

One of the main driver of forest degradation in Nepal is the demand for firewood, used as a source of cooking and heating energy. Thus, in 2010, rural households in the Hills and the Mountains collected on average five cubic meter of firewood per year.¹⁸ Using the large scale household surveys NLSSII and NLSSIII data, we analyse below household energy choices by estimating the following equations:¹⁹

$$Y_{hvt} = \alpha CFUG_{vt} + \mathbf{X}_{vt}\beta + \mathbf{W}_{ht}\gamma + \delta_d + \tau_t + \varepsilon_{hvt} \quad (4)$$

where the dependent variable Y stands for the number of bharis of firewood collected by household h in year t in village v or the number of hours it takes to collect one bhari of firewood. The main explanatory variable is $CFUG$, the share of the village area managed by CFUG in year t . As above, we also include a large set of village level control variables \mathbf{X} , household level controls \mathbf{W} , belt-zone fixed effects δ and survey wave fixed effects τ . ε_{hvt} is the idiosyncratic component. The coefficient of interest is α . It indicates how cross sectional variations in the share of VDC area managed by CFUG at time t are related to changes in the dependent variable Y . Given the previous discussion about the endogenous placement of CFUG, this is not a causal estimate, a point that we discuss below. Taking into account the forest regeneration process driven by CFUGs, we also investigate the possibility that the effects of the CFUGs vary with time by distinguishing between new and old (more than 15 years) CFUGs.

The first two columns of table 7 report the estimations for the firewood collection time. The presence of community forest user groups is typically correlated to larger collection times. As column (1) indicates, a 10% increase in CFUG area

¹⁸Households report 79 bharis (headload) of firewood. We converted this by assuming that one bhari weights 30kg and 500 kg of wood correspond to one cubic meter.

¹⁹We could not use NLSSI data due to lack of information on some important control variables, such as rainfall or temperature.

is associated with an increase of 0.12 hours in collection time. This correlation vanishes for older CFUG (more than 15 years of age) as the sum of the coefficient estimated for current CFUG coverage and this coverage 15 years ago is insignificant and close to zero (column (2)).²⁰ These results indicate that, if anything, CFUGs tend to initially restrict the access to forest and households have to rely on forests located further away for firewood. Once the forest regenerates, these restrictions are gradually relaxed.

[Insert Table 7 here]

Quite surprisingly, the presence of CFUGs does not, on average, correlates with the amounts of firewood collected (column 3 of table 7. This is probably an artefact as, when we consider separately young and old CFUG, we find that young CFUG are associated with significantly lower levels of firewood collection while older CFUG have, if anything, a net positive effect on collections. This is consistent with our previous results on forest expansion but also with repeated stories from our field interviews claiming that, after a restriction period, improved forest management raises biomass production and provide more forest products to villagers. In the last two columns, we introduce firewood collection time as an endogenous control, as it represents the main direct cost associated with household firewood collection. It reduces the point estimates of the CFUGs coefficients, but only partly. This suggests that CFUGs may also affect the demand for energy in other ways than through variations in collection time, for instance by collecting collection fees, banning access to fragile areas or restricting collections to specific periods and dry wood.

Finally, we also investigate how CFUGs coverage in a village affect other fuel expenditures. Table 8 indicates a (weakly significant) positive correlation between CFUG coverage and the amounts spent on fuel by the households. Thus, in villages with 10% more CFUG coverage than the other villages, households spend 158NPR more on fuel (the average fuel expenditures per household are equal to 2100NPR). This relation seems to be essentially driven by firewood collection times.

²⁰Since 7% of rural households do not collect firewood, they don't report collection time. As a robustness check, we also estimate a village level regression by taking the village median collection time as the dependent variable. We report estimation results in table A3 with similar results.

[Insert Table 8 here]

This last set of results is based on a cross-section of households and suffers from potential endogeneity issues in the creation of a CFUG. As discussed previously, there are reasons to believe that the forests handed over to the communities were mostly degraded forests, lying close to market centres and under large pressure by the users. Under this argument, older CFUGs correspond to places where households enjoy better access to markets and alternative energy sources. To reduce these potential biases, we control for a large set of observable factors that can influence both forest conditions, CFUG creation and household behaviour, such as the distance between the village and a paved road, the distance to district headquarters or population density and household level characteristics such as access to land, livestock ownership, non-farm business assets or the number of migrants in the households. But this may not be sufficient. In particular, the selection problem should be more severe around the creation date of the CFUG than several years after their creation. In this sense, the fact that more recent CFUGs are associated to lower collection levels and higher collection time while older CFUG, created closer to the district center, are associated to higher collections and lower collection times is reassuring, confirming CFUG effectiveness as time passes. This is very much in line with related findings on the Indian Himalayas (Baland et al., 2010).²¹

5 Discussion and conclusion

In this paper, we first assess the positive contribution of community forestry in the Hills and Mountains of Nepal on forest regeneration, using an instrumental variable approach in the spirit of a program roll-out. We then investigate the potential mechanisms driving this increase and show that CFUGs played a role both on demand, for instance by increasing the costs of firewood collection in the short run, and supply, by increasing the size of the forests and changing their composition. A few remarks are in order that help to qualify the interpretation of

²¹We did try to instrument the share of the area managed by CFUG, following the approach described in equation (2) but the instrument, even if statistically significant, is not strong enough given the smaller number of villages in the household level approach.

our results. First, the presence of CFUGs is measured at the “village” level and not at the plot level. We therefore measure the average effect of CFUG at the village level, both on forest plots that are managed by CFUGs but also on nearby areas in the village. In terms of interpretation, the effect that we highlight is therefore net of spillovers across plots within the same village. If the CFUGs protect the forests under their control, increasing tree cover, but induce increased pressure and forests degradation in nearby plots, we actually measure the net weighted average of these two effects.

Environmental awareness or better access to new sources of energy may also reduce the pressure on overall forest resources. Moreover, the development of community forestry may also encourage the expansion of trees on other plots of land, as villagers may start planting trees on their private plots to compensate for the reduced access to firewood and fodder from common land. All these are also accounted for in the average effect we estimate at the village level. Whatever the precise mechanisms, we believe that the net effect we highlight is crucial for policy makers as it shows that the program is on average positive for forests in Nepal. Note however that we could not exclude negative spillovers on neighbouring villages in which CFUGs are absent. Given the size of the villages, and the large number of controls we used, we believe however that this potential bias is arguably negligible. Using a different methodology, we intend to evaluate the importance of these within and across village spillovers in future research.

Also, our study stresses the importance of distinguishing between short and long run effects in the context of natural resources, such as forests, as they take time to regenerate. Shortly after their creation, CFUGs typically restrict access to forest resources, such as firewood or timber. When effective, these restrictions lead to larger, richer and more dense forests, allowing in the long run better harvests than at the time of the CFUG creation. Thus, in Nepal, we show that the amounts of firewood collected in villages with older CFUGs are similar to those in villages without CFUGs and larger than the average collection levels in places with young CFUGs. Our field visits also indicate that several old CFUGs are now able to also supply timber in a sustainable manner and actually generate sizeable incomes. The long term success of CFUGs is however conditioned to the short-term ability to reduce the demand for forest products. In our context, where firewood is an

important driver of forest degradation (Baland et al., 2018), access to other sources of energy is of paramount importance. More broadly, this implies that the development of a community forestry program with the goal of restoring forests should go hand in hand with a proper understanding of their main use. Well-designed policies should therefore provide temporary solutions to alleviate the burden of forest conservation on the regular users of the forests. These solutions can then be lifted when the forest is again dense enough to provide ecosystem services in a sustainable manner.

By focussing on the average benefits of community forestry at the local level, we could not investigate the distribution of its costs and benefits across villagers. Thus, more ecosystem services also mean that the population of wild animals increases, causing crop damages in nearby cultivated plot or killing poultry and livestock as several villagers told us during field work and as mentioned by Baral et al. (2021) around some community forest in the Mid-Hills of Nepal.²² The distribution of forests benefits also changes as a consequence of community forests. For instance, a typical claim of women is that men leading CFUGs focus on pine trees that can be sold as timber whereas they would prefer more broadleaf trees as a source of firewood and fodder, a task traditionally devoted to women (see in particular Agarwal (2010), Leone (2019) and Bocci and Mishra (2021)).²³

Community forestry in Nepal is a game changer at the local level. This institutional change empowers local communities to restore degraded forests and possibly escape a poverty-environment vicious circle (Dasgupta and Mäler, 1995). At the global level, community forestry in Nepal increases carbon sequestration and contributes to the mitigation of global warming. Our study probably overestimates this contribution as some of the reductions in firewood used is partly compensated by the use of other energy sources. Clearly, the development of biogas is beneficial from climate. However, when these alternative energy sources come from the market, under the form of charcoal, firewood collected further away, LPG or kerosene, the pure local effect of CFUGs on forest restoration overestimates their contribution to climate change mitigation.

²²See Gulati et al. (2021) for a broader discussion of the costs of human wildlife conflict around protected areas in nearby India.

²³The change in the composition of forests may also have adverse effect in terms of biodiversity by favouring species that prefer coniferous forests over broadleaf forests or bocage like landscapes.

Acknowledgements

The authors are grateful to Julia Girard and Camille Roegiers for their valuable research assistance. The authors benefited from the suggestions and advice from Olivia Aubriot, Mani Nepal, E. Somanathan, Jean-Philippe Platteau and Joelle Smadja. This work also benefited from numerous constructive comments during the EAERE, the EDI and the CECFEE annual conference as well as during workshops and seminar at RWI Essen, ECOPUB Paris, PSE Paris, TSE toulouse, BETA Nancy, UPPA Pau and CERDI Clermont-Ferrand. This research received the support of the EDI research program and the ANR GOLFOR-DEEPN (ANR-18-CE03-0005). François Libois also acknowledges the support of a CEPREMAP seed funding and of a French government subsidy managed by the Agence Nationale de la Recherche under the framework of the Investissements d'avenir programme referenced ANR-17-EURE-001. The usual disclaimer applies and all mistakes remain author's responsibility. There is no conflict of interest to disclose

References


- Abrams, M. and Crippen, R. (2011). Aster gdem v3. Technical report, Japan's Ministry of Economy, Trade, and Industry (METI) and National Aeronautics and Space Administration (NASA).
- Agarwal, B. (2010). *Gender and Green Governance: The Political Economy of Women's Presence Within and Beyond Community Forestry*. Oxford University Press.
- Aitken, R. (1977). Wilderness areas in scotland. Technical report.
- Baland, J.-M., Bardhan, P., Das, S., and Mookherjee, D. (2010). Forests to the people: Decentralization and forest degradation in the Indian Himalayas. *World Development*, 38(11):1642 – 1656.
- Baland, J.-M., Libois, F., and Mookherjee, D. (2018). Forest degradation and economic growth in nepal, 2003–2010. *Journal of the Association of Environmental and Resource Economists*, 5(2):401–439.
- Baral, K., Sharma, H. P., Kunwar, R., Morley, C., Aryal, A., Rimal, B., and Ji, W. (2021). Human wildlife conflict and impacts on livelihood: A study in community forestry system in mid-hills of nepal. *Sustainability*, 13(23).
- Bluffstone, R. A., Somanathan, E., Jha, P., Luintel, H., Bista, R., Toman, M., Paudel, N., and Adhikari, B. (2018). Does collective action sequester carbon? evidence from the nepal community forestry program. *World Development*, 101:133–141.
- Bocci, C. and Mishra, K. (2021). Forest power: The impact of community forest management on female empowerment. *Ecological Economics*, 187:107105.
- Bojinski, S., Verstraete, M., Peterson, T. C., Richter, C., Simmons, A., and Zemp, M. (2014). The concept of essential climate variables in support of climate research, applications, and policy. *Bulletin of the American Meteorological Society*, 95(9):1431 – 1443.

- Bowler, D. E., Buyung-Ali, L. M., Healey, J. R., Jones, J. P., Knight, T. M., and Pullin, A. S. (2012). Does community forest management provide global environmental benefits and improve local welfare? *Frontiers in Ecology and the Environment*, 10(1):29–36.
- Bréda, N. (2008). Leaf area index. In Jorgensen, S. E. and Fath, B. D., editors, *Encyclopedia of Ecology*, pages 2148–2154. Academic Press, Oxford.
- CBS (2011). Nepal Living Standards Survey Household 2010-2011. Statistical Report Vol. 1, Nepal Central Bureau of Statistics.
- Dasgupta, P. and Mäler, K.-G. (1995). Chapter 39 poverty, institutions, and the environmental resource-base. In *Handbook of Development Economics*, volume 3 of *Handbook of Development Economics*, pages 2371–2463. Elsevier.
- de Chaisemartin, C. and D’Haultfoeuille, X. (2020). Two-way fixed effects estimators with heterogeneous treatment effects. *American Economic Review*, 110(9):2964–96.
- Desbureaux, S. (2016). Common resources management and the “dark side” of collective action: an impact evaluation for madagascar’s forests.
- Edmonds, E. V. (2002). Government-initiated community resource management and local resource extraction from Nepal’s forests. *Journal of Development Economics*, 68(1):89–115.
- Friedl, M., McIver, D., Hodges, J., Zhang, X., Muchoney, D., Strahler, A., Woodcock, C., Gopal, S., Schneider, A., Cooper, A., Baccini, A., Gao, F., and Schaaf, C. (2002). Global land cover mapping from modis: algorithms and early results. *Remote Sensing of Environment*, 83(1):287–302. The Moderate Resolution Imaging Spectroradiometer (MODIS): a new generation of Land Surface Monitoring.
- Friedl, M. A., Sulla-Menashe, D., Tan, B., Schneider, A., Ramankutty, N., Sibley, A., and Huang, X. (2010). Modis collection 5 global land cover: Algorithm refinements and characterization of new datasets. *Remote Sensing of Environment*, 114(1):168–182.

- Fujisada, H., Urai, M., and Iwasaki, A. (2011). Advanced methodology for aster dem generation. *IEEE Transactions on Geoscience and Remote Sensing*, 49(12):5080–5091.
- Gulati, S., Karanth, K. K., Le, N. A., and Noack, F. (2021). Human casualties are the dominant cost of human–wildlife conflict in india. *Proceedings of the National Academy of Sciences*, 118(8).
- Hall, D. K. and Riggs, G. A. (2021). Modis/terra snow cover 8-day l3 global 500m sin grid. *MODIS/Terra Snow Cover 8-Day L3 Global 500m SIN Grid, Version 61*.
- Hatlebakk, M. (2007). LSMS data quality in Maoist influenced areas of Nepal. CMI Working Papers 6, CMI (Chr. Michelsen Institute), Bergen, Norway.
- ICIMOD (2014a). Land cover of nepal 2000.
- ICIMOD (2014b). Land cover of nepal 2010.
- INSEC (2009). *Electronic data files*. Informal Service Service Center, Kathmandu.
- Joshi, M. and Pyakurel, S. R. (2015). Individual–level data on the victims of nepal’s civil war, 1996–2006: A new data set. *International Interactions*, 41(3):601–619.
- Langmuir, E. (2013). *Mountaincraft and Leadership*. Mountain Leader Training England.
- Leone, M. (2019). Women as decision makers in community forest management: Evidence from nepal. *Journal of Development Economics*, 138:180–191.
- Libois, F. (2016). Households in times of war: Adaptation strategies during the Nepal civil war. Technical report, CRED, University of Namur.
- Libois, F. (2021). Success and failure of communities managing natural resources: Static and dynamic inefficiencies. Technical report, Paris School of Economics and INRAE.

- Niraula, R. R., Gilani, H., Pokharel, B. K., and Qamer, F. M. (2013). Measuring impacts of community forestry program through repeat photography and satellite remote sensing in the dolakha district of nepal. *Journal of Environmental Management*, 126:20–29.
- Oldekop, J. A., Sims, K. R. E., Karna, B. K., Whittingham, M. J., and Agrawal, A. (2019). Reductions in deforestation and poverty from decentralized forest management in nepal. *Nature Sustainability*, 2(5):421–428.
- Somanathan, E., Prabhakar, R., and Mehta, B. S. (2009). Decentralization for cost-effective conservation. *Proceedings of the National Academy of Sciences of the United States of America*, 106(11):4143–4147.
- Sukhdev, P. and Nuttall, N. (2010). Green economy: A brief for policymakers on the green economy and millennium development goals.
- Sulla-Menashe, D. and Friedl, M. A. (2018). User guide to collection 6 modis land cover (mcd12q1 and mcd12c1) product. Technical report, Boston University.
- TRMM (2011). Trmm (tmpa/3b43) rainfall estimate l3 1 month 0.25 degree x 0.25 degree v7.
- Verger, A., Baret, F., and Weiss, M. (2014). Near real-time vegetation monitoring at global scale. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing*, 7(8):3473–3481.
- Wan, Z., Hook, S., and Hulley, G. (2015). Mod11a2 modis/terra land surface temperature/emissivity 8-day l3 global 1km sin grid v006 [data set]. *NASA EOSDIS Land Processes DAAC*.

Figure 1: Leaf Area Index and CFUG in the Hills and the Mountains of Nepal



Figures

Figure 2: Community forestry expansion in Nepal




Figure 3: Forest cover and Community Forest User Group through time




Figure 4: Short and long term effects of CFUG creation on Leaf Area Index




Figure 5: Estimation based on de Chaisemartin and D'Haultfoeuille (2020)), VDC share has been re-categorized in 6 categories, 0 for VDC without CFUG, 1 for >0-20%, 2 fro 20-40%, etc... The standard errors computed are based on 500 bootstrap replicates.

Tables

Table 1: Descriptive statistics: VDC level variables, Hills and Mountains 2001-2013

Variable	2001			2013			all years		
	Median	Mean	Std. dev.	Median	Mean	Std. dev.	Median	Mean	Std. dev.
November Leaf Area Index	1.14	1.05	.69	1.43	1.33	.92	1.24	1.15	.78
December Leaf Area Index	.82	.77	.5	.97	.88	.6	.91	.85	.57
Share VDC managed by CFUG	.02	.08	.13	.05	.12	.16	.03	.11	.15
VDC area (ha.)	2227.08	4430.01	9141.19	2227.08	4430.01	9141.19	2227.08	4430.01	9139.56
Share forested in 1950	.45	.46	.32	.45	.46	.32	.45	.46	.32
Walking distance to HQ (hrs.)	8.86	10.63	6.48	8.86	10.63	6.48	8.86	10.63	6.48
FUG years in district	10	9.85	1.62	22	21.85	1.62	16	16.35	3.81
Total precipitation	1732.26	1665.9	539.5	1710.98	1570.47	488.67	1519.74	1469.14	526.68
Average snow cover	.01	.02	.02	.01	.02	.02	.01	.02	.02
Growing degree days	2996.74	2914.47	2066.47	2741.87	2725.5	1945.53	2851.22	2810.59	1960.65
Conflict related casualties	3	7.16	12.61	0	0	0	0	6.02	17.56
Population density	.48	1.04	3.66	.56	1.18	5.98	.53	1.11	4.82
Biogas per household	0	.01	.02	0	.02	.07	0	.02	.05
Observations		2578			2578			30936	

Descriptive statistics for the panel of VDC in the Hills and Mountains

Observations weighted by VDC area (at the exception of VDC area)

Table 2: Descriptive statistics: household level variables

Variable	2003		2010		full sample	
	Mean	Median	Mean	Median	Mean	Median
Wood	85.94 (55.4)	72	78.91 (61.68)	60	81.78 (59.29)	70
Collection time	3.5 (1.69)	3	3.91 (1.9)	4	3.74 (1.83)	3.5
Fuel expenditures	1387.19 (2622.16)	813.45	2578.22 (4554.36)	884.47	2091.56 (3926.27)	845.98
% of Vil. area in FUG	.14 (.14)	.1	.2 (.19)	.14	.18 (.18)	.13
% of Vil. area in FUG 15 years ago	0 (0)	0	.05 (.1)	.02	.03 (.08)	0
Years since 1st CFUG in district	12.54 (1.66)	12	19.42 (1.58)	19	16.61 (3.75)	18
Walking time to district HQ	5.96 (4.01)	5.5	6.07 (3.76)	5.43	6.03 (3.86)	5.45
Forest cover in 1950	.37 (.33)	.25	.42 (.31)	.44	.4 (.32)	.35
Observations	1474		2116		3578	

Descriptive statistics for the second and third repeated cross-sections of NLSS in rural villages.

All monetary values expressed in NPR2010

Standard errors in parentheses

Table 3: Change in November Leaf Area index as a function of CFUG expansion

	Panel F.E. (1)	First stage (2)	Panel F.E. + IV (3)	Panel F.E. (4)	First stage (5)	Panel F.E. + IV (6)
Share FUG in VDC	0.448*** [7.36]		4.594*** [4.97]	0.366*** [7.14]		5.281*** [4.29]
Proximity Hq × FUG years in district		0.00546*** [4.58]			0.00449*** [4.19]	
Population density				-0.00356** [-2.34]	-0.00115*** [-2.75]	-0.00254* [-1.89]
Biogas per household				0.878*** [5.84]	0.253*** [3.64]	-0.507 [-1.18]
Years since FUG in district	0.0221*** [6.50]	-0.00703*** [-3.23]	0.169*** [7.59]	0.0206*** [6.26]	-0.00764*** [-3.53]	0.178*** [7.38]
Forest in 1950 × FUG years in district	-0.000184 [-0.05]	0.00496*** [3.31]	-0.0197*** [-3.38]	0.00129 [0.43]	0.00547*** [3.97]	-0.0248*** [-3.35]
VDC fixed effects	Yes	Yes	Yes	Yes	Yes	Yes
Year fixed effects	Yes	Yes	Yes	Yes	Yes	Yes
Environment controls	Yes	Yes	Yes	Yes	Yes	Yes
Observations	136252129	136252129	136252129	131392040	131392040	131392040
Observations (in VDC)	2552x13	2552x13	2552x13	2471x13	2471x13	2471x13
Mean LAI in 2013	1.33					
Mean FUG share in 2013	.12					

Regressions are weighted by VDC area. Environment controls include rainfall, snow cover, growing degree days and conflict related casualties. We derive population data from the 2011 and 2011 population census and interpolate figures.

Standard errors clustered at the district level

t-statistics in brackets, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table 4: Land use change as a function of CFUG expansion in the Hills and Mountains

	Panel F.E.			1st stage		Panel F.E + IV	
	(Forest) (1)	(Needle) (2)	(Mixed) (3)	(Share FUG) (4)	(Forest) (5)	(Needle) (6)	(Mixed) (7)
Share FUG in VDC	0.0764*** [4.69]	0.00821** [2.17]	0.0661*** [4.49]		0.583*** [3.14]	0.0686* [1.84]	0.405** [2.29]
Proximity Hq × FUG years in district				0.00547*** [4.59]			
Years since FUG in district	0.00138*** [3.12]	-0.000121* [-1.78]	0.00174*** [4.23]	0.000249 [0.53]	0.000781* [1.78]	-0.000193** [-2.01]	0.00133*** [3.17]
Forest in 1950 × FUG years in district	-0.00258*** [-4.00]	0.0000510 [0.27]	-0.00164** [-2.60]	0.00497*** [3.32]	-0.00497*** [-4.63]	-0.000234 [-0.78]	-0.00324*** [-3.61]
VDC fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Environmental controls	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Observations (in ha)	139495824	139495824	139495824	139495824	139495824	139495824	139495824
Observations (VDCxyear)	2552x13	2552x13	2552x13	2552x13	2552x13	2552x13	2552x13

Regressions are weighted by VDC area. Environment controls include rainfall, snow cover, growing degree days and conflict related casualties.

Standard errors clustered at the district level, regression weighted by VDC area

t-statistics in brackets, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table 5: Land use change as a function of CFUG expansion in the Hills and Mountains

	Panel F.E.			1st stage		Panel F.E + IV	
	(Broadleaf) (1)	(Crop) (2)	(Shrub) (3)	(Share FUG) (4)	(Broadleaf) (5)	(Crop) (6)	(Shrub) (7)
Share FUG in VDC	0.00203 [0.26]	-0.0172*** [-2.95]	-0.0735*** [-4.32]		0.109 [1.61]	-0.390*** [-3.22]	-0.494*** [-3.02]
Proximity Hq × FUG years in district				0.00547*** [4.59]			
Years since FUG in district	-0.000230* [-1.83]	-0.00115*** [-4.23]	0.0000642 [0.15]	0.000249 [0.53]	-0.000358** [-2.40]	-0.000702*** [-2.78]	0.000565 [1.38]
Forest in 1950 × FUG years in district	-0.000990*** [-3.60]	0.00136*** [3.25]	0.00170** [2.18]	0.00497*** [3.32]	-0.00149*** [-3.45]	0.00312*** [3.66]	0.00368*** [3.21]
VDC fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Year fixed effects	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Environmental controls	Yes	Yes	Yes	Yes	Yes	Yes	Yes
Observations (in ha)	139495824	139495824	139495824	139495824	139495824	139495824	139495824
Observations (VDCxyear)	2552x13	2552x13	2552x13	2552x13	2552x13	2552x13	2552x13

Regressions are weighted by VDC area. Environment controls include rainfall, snow cover, growing degree days and conflict related casualties.

Standard errors clustered at the district level, regression weighted by VDC area

t-statistics in brackets, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table 6: Biogas adoption in the Hills and Mountains

	Panel F.E.		1st stage	Panel F.E + IV	
	(Biogas units) (1)	(Biogas units per hh.) (2)	(Share FUG) (3)	(Biogas units) (4)	(Biogas units per hh.) (5)
Share FUG in VDC	180.9*** [4.26]	0.0743*** [3.95]		4074.0*** [2.93]	0.848*** [3.49]
Years since FUG in district	2.229* [1.94]	0.00142*** [2.83]	0.000249 [0.53]	-2.413 [-1.22]	0.000499 [1.16]
Forest in 1950 × FUG years in district	0.627 [0.17]	-0.000796 [-0.71]	0.00497*** [3.32]	-17.75** [-2.06]	-0.00445*** [-3.25]
Proximity Hq × FUG years in district			0.00547*** [4.59]		
VDC fixed effects	Yes	Yes	Yes	Yes	Yes
Year fixed effects	Yes	Yes	Yes	Yes	Yes
Environmental controls	Yes	Yes	Yes	Yes	Yes
Observations (in ha)	139495824	139495824	139495824	139495824	139495824
Observations (VDCxyear)	2552x13	2552x13	2552x13	2552x13	2552x13

Regressions are weighted by VDC area. Environment controls include rainfall, snow cover, growing degree days and conflict related casualties.

Standard errors clustered at the district level, regression weighted by VDC area

t-statistics in brackets, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table 7: Firewood collection

	collection time (hrs)		Firewood collection (bhari)			
	(1)	(2)	(3)	(4)	(5)	(6)
% of Vil. area in FUG	1.218*** (2.82)	1.471*** (3.18)	-15.29 (-1.35)	-27.94** (-2.43)	-9.193 (-0.83)	-20.94* (-1.83)
% of Vil. area in FUG 15 years ago		-1.160 (-1.09)		65.47** (2.51)		58.05** (2.44)
Med. collection time					-4.244*** (-2.99)	-3.876*** (-2.83)
Years since 1st CFUG in district	-0.0159 (-0.37)	-0.0113 (-0.27)	1.988 (1.40)	1.736 (1.24)	1.837 (1.33)	1.627 (1.19)
Proximity to district HQ	0.0334 (1.44)	0.0331 (1.44)	1.225** (2.32)	1.245** (2.38)	1.311** (2.55)	1.322** (2.58)
Forest cover in 1950	-0.00520 (-0.02)	-0.0247 (-0.09)	3.988 (0.57)	4.834 (0.67)	4.987 (0.73)	5.651 (0.81)
Household assets	Yes	Yes	Yes	Yes	Yes	Yes
Year fixed-effect	Yes	Yes	Yes	Yes	Yes	Yes
Belt-Zone fixed-effects	Yes	Yes	Yes	Yes	Yes	Yes
Village controls	Yes	Yes	Yes	Yes	Yes	Yes
Observations	3332	3332	3578	3578	3578	3578

Village controls include distance to paved road, war casualties, median elevation and standard deviation, snow cover, rainfall, growing degree days and cooling degree days

Standard errors clustered at the village level – t -statistics in parentheses, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table 8: Fuel expenditures

	Fuel expenditures (NPR)			
	(1)	(2)	(3)	(4)
% of Vil. area in FUG	1581.6*	1756.1*	1044.8	1083.3
	(1.91)	(1.84)	(1.45)	(1.31)
% of Vil. area in FUG, 15 years ago		-902.7		-190.2
		(-0.44)		(-0.11)
Med. collection time			373.6***	372.4***
			(2.76)	(2.77)
Years since 1st CFUG in district	-277.8**	-274.3**	-264.6**	-263.9**
	(-2.54)	(-2.51)	(-2.40)	(-2.39)
Proximity to district HQ	-86.70*	-86.98*	-94.32**	-94.36**
	(-1.83)	(-1.84)	(-2.03)	(-2.03)
Forest cover in 1950	-795.1*	-806.8*	-883.1**	-885.3**
	(-1.82)	(-1.83)	(-1.99)	(-1.99)
Household assets	Yes	Yes	Yes	Yes
Year fixed-effect	Yes	Yes	Yes	Yes
Belt-Zone fixed-effects	Yes	Yes	Yes	Yes
Village controls	Yes	Yes	Yes	Yes
Observations	3578	3578	3578	3578

Village controls include distance to paved road, war casualties, median elevation and standard deviation, snow cover, rainfall, growing degree days and cooling degree days

Standard errors clustered at the village level – t -statistics in parentheses, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Figure A1: CFUG creation year



A Appendix

Figure A2: Area managed by community forest user groups in 1993
 Share of municipality area managed by CFUG
 1993




Figure A3: Area managed by community forest user groups in 1996
 Share of municipality area managed by CFUG
 1996




Figure A4: Area managed by community forest user groups in 2016




Figure A5: CFUG main forest type




Table A1: Descriptive statistics: household level variables (continued)

Variable	2003		2010		full sample	
	Mean	Median	Mean	Median	Mean	Median
Big livestock	3.56 (2.91)	3	3.15 (2.56)	3	3.32 (2.71)	3
Land owned, ha	.69 (.76)	.49	.61 (.66)	.43	.64 (.71)	.46
Household size	5.03 (2.24)	5	4.79 (2.16)	5	4.89 (2.2)	5
Prop. female	.35 (.19)	.33	.37 (.19)	.33	.36 (.19)	.33
Prop. children	.39 (.24)	.4	.37 (.24)	.4	.38 (.24)	.4
Avg. education	2.43 (2.7)	1.67	3.16 (2.98)	2.67	2.86 (2.89)	2.33
= 1 if NFBus	.22 (.41)	0	.28 (.45)	0	.26 (.44)	0
# Migrants	.4 (.67)	0	.8 (.97)	1	.64 (.88)	0
Med. time to road	10.17 (13.44)	5	5.68 (7.61)	2.5	7.52 (10.62)	3.13
# killings 20km ar.	79.40 (64.56)	56	151.18 (97.46)	126	121.85 (92.54)	101
Vil. elevation: mean	1426.39 (738.68)	1336	1478.55 (782.89)	1332	1457.24 (765.46)	1332
Vil. elevation: std. dev.	329.38 (206.06)	290.03	334.56 (208.08)	301.69	332.44 (207.25)	296.55
Vil. snow cover	0 (.01)	0	0 (.01)	0	0 (.01)	0
Rainfall z-score	.61 (.64)	.75	-.93 (.65)	-.91	-.3 (.99)	-.45
Monsoon GDD	1242.86 (364.87)	1361.57	1137.35 (419.61)	1257.03	1180.46 (401.47)	1310.46
Cooling Degree Days	166.98 (496.01)	16.14	166.91 (495.14)	9.32	166.94 (495.43)	14.74
Observations	1474		2116		3578	

Descriptive statistics for the second and third repeated cross-sections of NLSS in rural villages.

All monetary values expressed in NPR2010

Standard errors in parentheses

Table A2: Change in December Leaf Area index as a function of CFUG expansion

	Panel F.E. (1)	First stage (2)	Panel F.E. + IV (3)	Panel F.E. (4)	First stage (5)	Panel F.E. + IV (6)
Share FUG in VDC	0.448*** [7.36]		4.594*** [4.97]	0.366*** [7.14]		5.281*** [4.29]
Proximity Hq × FUG years in district		0.00546*** [4.58]			0.00449*** [4.19]	
Population density				-0.00356** [-2.34]	-0.00115*** [-2.75]	-0.00254* [-1.89]
Biogas per household				0.878*** [5.84]	0.253*** [3.64]	-0.507 [-1.18]
Years since FUG in district	0.0221*** [6.50]	-0.00703*** [-3.23]	0.169*** [7.59]	0.0206*** [6.26]	-0.00764*** [-3.53]	0.178*** [7.38]
Forest in 1950 × FUG years in district	-0.000184 [-0.05]	0.00496*** [3.31]	-0.0197*** [-3.38]	0.00129 [0.43]	0.00547*** [3.97]	-0.0248*** [-3.35]
VDC fixed effects	Yes	Yes	Yes	Yes	Yes	Yes
Year fixed effects	Yes	Yes	Yes	Yes	Yes	Yes
Environment controls	Yes	Yes	Yes	Yes	Yes	Yes
Observations	136252129	136252129	136252129	131392040	131392040	131392040
Observations (in VDC)	2552x13	2552x13	2552x13	2471x13	2471x13	2471x13
Mean LAI in 2013	.88					
Mean FUG share in 2013	.12					

Regressions are weighted by VDC area. Environment controls include rainfall, snow cover, growing degree days and conflict related casualties. We derive population data from the 2011 and 2011 population census and interpolate figures.

Standard errors clustered at the district level

t-statistics in brackets, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Table A3: Village median collection time

	Collection time (hr. per bhari)			
	(1)	(2)	(3)	(4)
% of Vil. area in FUG	1.443*** (3.18)	1.819*** (3.18)	1.677*** (2.82)	1.658*** (2.74)
% of Vil. area in FUG, 15 years ago		-1.940* (-1.89)	-1.740 (-1.63)	-1.745* (-1.71)
Current leaf area index				-0.273 (-1.44)
Years since FUG in district	-0.0390 (-0.85)	-0.0314 (-0.70)	-0.0270 (-0.45)	-0.0247 (-0.43)
Forest cover in 1950	0.247 (0.64)	0.221 (0.57)	0.340 (0.86)	0.400 (1.01)
Proximity to district HQ	0.0209	0.0202	0.0215	0.0192
Year fixed-effect	Yes	Yes	Yes	Yes
Belt-Zone fixed-effects	Yes	Yes	Yes	Yes
Village controls	Yes	Yes	Yes	Yes
Village asset density	Yes	Yes	Yes	Yes
Observations	300	300	300	300

Standard errors clustered at the district level – t -statistics in parentheses, * $p < 0.1$, ** $p < 0.05$, *** $p < 0.01$

Village level controls include median access time to road, village median altitude and

altitude standard deviation, number of people killed in the 20km around the village in the previous year,

as well as previous year snow cover, rainfall deviation, cooling degree days and monsoon growing degree days.